

Irradiation Effect of Low-Level Laser on Functional Outcome in Children with Bell's Palsy

Enas Elsayed*, PTD and Khaled A. Mamdouh**, PTD

*Department of Physical Therapy for Neuromuscular Disorders and its Surgery, Faculty of Physical Therapy, Cairo University.

**Department of physical therapy, Faculty of applied medical sciences, King Abdul-Aziz University

ABSTRACT

Background: There are many evidences that low-level laser therapy (LLLT) may stimulate nerve regeneration. **Objective:** The aim of the study was to evaluate the effectiveness of LLLT in enhancing functional recovery in children with Bell's palsy (BP). **Subjects:** Twenty patients with BP of axonotemetic type were included (thirteen males and seven females). They ranged in age between 12 and 15 years old. They were randomly divided into two groups of equal number, study and control. **Methods:** The severity of BP was assessed by "House-Brackmann grading scale". Also, facial symmetry was measured at rest by using tape measurement. Both groups received physical therapy program. Additionally, the control group received placebo laser while the study group received continuous infrared (IR) laser 830 nm, 100mW with dose of 6.0 J/cm², at five points overlying the main facial trunk and four motor branches of the facial nerve. The study continued for four weeks, three times / week, every session continued for 60 minutes. **Results:** The results revealed no significant difference between the study and the control groups after treatment, neither in House-Brackmann grades nor in facial symmetry measurement. **Conclusion:** It can be concluded that infrared laser, with the parameters used in the current study, was not effective in improving functional outcome in BP.

Keywords: House-Brackmann grading scale, Infrared laser, Bell's palsy.

INTRODUCTION

Peripheral injury of the facial nerve is a frequent disorder. It is a stressful situation for the patient and is functionally hazardous for the cornea. The idiopathic facial paralysis or Bell's palsy (BP) is the most common type¹⁶. It is an acute, lower motor neuron paralysis of the face, usually unilateral, related to inflammation and swelling of facial nerve within the facial canal or at the stylomastoid foramen⁸. The anatomy of the facial nerve is rather complex for a cranial nerve, with a long intracranial course, in which the nerve takes three bends or genu⁶.

The exact causes of BP are unknown. However, it is thought that viral infection of

the nerve is the most likely cause. Herpes simplex virus is considered the most likely viral cause^{10,12,13}. Moreover, the pathophysiology of BP remains unclear and the treatment also remains controversial¹⁶.

Patients with facial nerve disorders are often devastated due to the emotional and psychological impact of facial disfiguration as well as the subsequent physical limitations and difficulties associated with speaking, drinking, eating, and facial expression. Socialization and community participation is extraordinarily limited and difficult for many of those patients³.

Studies on BP are difficult to evaluate, bearing in mind the high spontaneous remission rate. Moreover, there are no

established guidelines for the treatment. The treatment strategy for BP is to accelerate recovery, to prevent further worsening of facial palsy in patients who are only mildly affected, and to avoid sequelae, such as synkinesis, contracture, and facial asymmetry, in more severe cases. Acceleration of damaged nerve regeneration is one important area of research. Low level laser therapy (LLLT) is an electrical modality that is used widely in the medical field. Individuals and families use laser instead of medication for variety of conditions ranging from headache, low back pain to insomnia. However, the swelling and the neuritis in the bony facial canal is believed to be influenced by LLLT in patients with facial palsy. This intra-bony region of the nerve is the major target of laser therapy in this indication. The suspected association with herpes virus also supports the use of laser therapy in such cases^{7,18}.

Aim of the study

The aim of the study was to evaluate the effectiveness of LLLT on the functional recovery in children with BP.

SUBJECTS, MATERIALS AND METHODS

Subjects:

Twenty patients with BP (thirteen males and seven females) were selected from King Fahd Armed Forces Hospital in Jeddah. Inclusion criteria: The patients' age ranged from 12 to 15 years (mean = 13 ± 1.1), the duration of illness ranged from 4 to 6 weeks, and the facial nerve injury was of axonotemetic type confirmed by electro-physiological examination. Exclusion criteria: Facial problems other than BP as wounds, burns, skin allergy conditions or diabetes mellitus. Patients were randomly assigned into two groups of equal number, study and control.

Instrumentation:

Laser instrument:

Enraf Nonius Endolaser, model 476, made in Argentina, AlGaAs diode laser with wavelength 760/830 nm. The beam delivery system is hand held probe, continuous/ pulsed, the output power ranges for 25-100 mw, and the dose ranges from 0.01-9.99 Joules. It has two accessory goggles for eye protection (Fig. 1).



Fig. (1): Laser instrument.

House-Brackmann grading scale: It was used to assess the severity of BP. It categorizes BP from normal (grade one) to total palsy (grade six) (table 1).

Table (1): House-Brackmann grading scale¹³.

GRADE	DESCRIPTION
1 (Normal)	Normal facial function in all nerve branches
2 (Slight)	Gross: Slight weakness in close inspection and slight synkinesis. At rest: Normal tone and symmetry. Motion: - Forehead: Good to moderate movement. - Eye: Complete closure with minimum effort. - Mouth: Slight asymmetry.
3 (Moderate)	Gross: Obvious but not disfiguring facial asymmetry. Synkinesis is noticeable but not severe. There may be hemi-facial spasm or contracture. At rest: Normal tone and symmetry. Motion: - Forehead: Slight to moderate movement. - Eye: Complete closure with effort. - Mouth: Slight weakness with maximum effort.
4 (Moderately severe)	Gross: Asymmetry is disfiguring and / or obvious facial weakness. At rest: Normal tone and symmetry. Motion: - Mouth: Asymmetrical with maximal effort.
5 (Severe)	Gross: Only slight, barely noticeable movement. At rest: Asymmetrical facial appearance. Motion: - Forehead: No movement. - Eye: Incomplete closure. - Mouth: Slight movement.
6 Total	No facial function.

Methods:

Evaluation:

All patients were assessed before and after treatment by the House-Brackmann grading scale of facial palsy. Facial symmetry was measured at rest by tape measurement recording the distances from the ear tragus to

the angle of the mouth. It was carried out for both sides of the face (affected and unaffected) in the study and control groups.

Treatment:**Control group:**

Patients received a physical therapy program which consisted of manual massage, exercising of the facial muscles performed in lying position at first, then in sitting. He/She was asked to look surprise then frown, close eyes firmly then open wide, smile, grin, say "o, e", whistle and blow. The patient received assistance at first then progressed to resistance. A mirror was useful to enable the patient to observe the muscle activity and compare between both sides of the face. The patient was also instructed to follow exercise program, twice a day, ten repetitions for each exercise with rest in between and in front of mirror¹⁷. In addition, they received placebo laser irradiation.

Study group:

Patients received the previously mentioned physical therapy program. In addition, they received LLLT. Before laser application, patient was prepared as following: They were instructed to remove facial hair before laser treatment. Then the area of treatment was cleaned well with alcohol. He/she assumed a comfortable position and was asked to wear the goggles.

Laser was applied on the main nerve trunk which lies 18.8 mm caudal to the tragus of the ear where it lies 23.6 mm deep to skin surface¹⁹. In addition, four motor branches of

the facial nerve (temporal, zygomatic, buccal and mandibular) were also irradiated about two centimeters from the nerve trunk in the direction of each branch. Laser parameters were continuous wavelength 830 nm, 100 mW, with dose of 6.0 J/cm² for 4 minutes for each point¹⁴. The treatment was continued for four weeks, three times / week; every session continued for 60 min.

Statistical analysis:

Data was presented as means and standard deviations. Paired and unpaired t-tests were used to analyze parametric data. Mann Whitney and Wilcoxon tests were used to analyze non-parametric data. The P-value was 0.05.

RESULTS**House-Brackmann grading scale:**

There was no significant difference between study and control groups regarding House-Brackmann grade mean values neither before (mean = 5.04 ± 0.21 and 5.08 ± 0.13, respectively) nor after treatment (mean = 4.87 ± 0.37 and 5 ± 0.14, respectively), (P = 0.7 and 0.8, respectively). Also, the results revealed no significant difference after treatment compared to that before treatment in both the study and control groups (P = 0.4 and 0.1 respectively) (table 2 and fig. 2).

Table (2): Comparison between study and control groups as regard House-Brackmann grade mean values before and after treatment.

House-Brackmann grades	Study group Mean ± SD	Control group Mean ± SD	P
Pre	5.04 ± 0.21	5.08 ± 0.13	0.7
Post	4.87 ± 0.37	5 ± 0.14	0.8
P	0.4	0.1	




Fig. (2): Mean values of House-Brackmann grades before and after treatment in the study and control groups.

Facial symmetry:

The results also revealed non significant difference between the study and control groups regarding facial symmetry measurement mean values neither before (mean = 9.6 ± 0.5 cm and 10.6 ± 0.6 cm respectively) nor after intervention (9.4 ± 0.55

and 10.3 ± 0.6 respectively) ($P = 0.18$ and 0.12 respectively). On the contrary, there was a significant decrease of measurement mean values within the study and control groups after treatment, compared to that before treatment ($P = 0.009$ and 0.05 respectively) (table 3 and fig. 3).

Table (3): Comparison between study and control groups as regard facial symmetry measurement before and after treatment.

House- Brackmann grading scale	Study group Mean \pm SD	Control group Mean \pm SD	P
Pre	9.6 ± 0.5	10.6 ± 0.6	0.18
Post	9.4 ± 0.55	10.3 ± 0.6	0.12
P	0.009 **	0.05 *	

* Significant

** Highly significant




Fig. (3): Mean values of facial symmetry measurement before and after treatment in the study and control groups.

The results also showed non significant difference in facial symmetry measurement between normal and affected sides of the face in the study group. The mean values of facial

symmetry measurement of the affected and unaffected sides were (9.4 ± 0.55 and 9.2 ± 0.55 respectively) after treatment ($P = 0.6$) (fig. 4).




Fig. (4): Mean values of facial symmetry measurement of the affected and unaffected sides after treatment in the study group.

DISCUSSION

Untreated BP leaves some patients with major facial dysfunctions and a reduced quality of life. The current study aimed at investigating the effect of IR laser on functional recovery in children with BP. Many evidences have shown that peripheral nerves regeneration may be accelerated by physical agents, such as electric power, magnetic fields, and ultrasound. Laser therapy has also been studied regarding a potential positive role in this particular area, with first investigations focusing on changes in nervous stimulus passage, with an electrophysiological demonstration of a reduced latency time, and increased conduction velocity in normal nerves in human beings^{2,11}.

The results of the current study revealed no significant difference between the study and control groups after laser irradiation regarding functional recovery according to House- Brackmann scale grades and facial symmetry measurements. On the other hand, a significant improvement in facial symmetry was found within the study and the control groups. This may be attributed to the effect of massage as well as the facial exercises that regained the face to normal or near normal alignment.

In accordance with the findings of the present work, Saitkulov et al.,¹⁵ studied the effect of IR laser irradiation on regeneration of nerve fibers in rat sciatic nerve. Starting from the first day after nerve crushing, the projection of proximal portion of the nerve was transcutaneously stimulated by 890 nm laser irradiation with total dose of 42 J/cm². On day 30 after crushing, there were no significant differences in the number of

myelinated fibers and mass of regenerating soleus muscle in the test and control groups.

Furthermore, Chen et al.,⁴ studied the effect of low-power pulsed laser irradiation on the regeneration of a 10-mm gap of rat sciatic nerve created between the proximal and distal nerve stumps. After eight weeks of recovery, pulsed laser-irradiated groups had significantly lower success rate of regeneration compared to sham-irradiated controls. They reported that deleterious effects have been demonstrated, such as lower regeneration percent rate translated by a less mature ultra structural organization, with smaller cross-sectional areas and smaller number of myelinated axons in irradiated nerves. They concluded that irradiation with pulsed laser may have suppressive effects on peripheral nerves regeneration.

On the contrary, Anders et al.,¹ examined quantitatively if transcutaneous LLL irradiation can affect the regeneration of the rat facial nerve which was crushed unilaterally in rats and irradiated daily with a laser directed to the area of the crush injury. It began on the day of the injury and was continued daily for 7, 8, or 9 days. Preliminary experiments determined the most effective wavelength, laser power, length of irradiation, and treatment schedule. The most effective laser parameters included daily irradiation with a helium-neon (HeNe) or argon pumped tunable dye laser (wavelength = 633 nm) with a power of 8.5 mW for 90 minutes (45.9 J, 162.4 J/cm²). The number of horseradish peroxidase (HRP) labeled neurons in the facial motor nucleus were used as an assay of the degree of regeneration. There was a statistically significant difference between the control and irradiated rats on day 9 postcrush. These data indicated that transcutaneous LLL irradiation

with the laser type and parameters involved in this study increased the rate of regeneration of rat facial nerve. It is evident that the difference between the previous results and the findings of the current work can be explained by the different laser type and parameters used.

Moreover, Endo et al.,⁵ tested IR laser irradiation in controlled crush injury of sciatic nerve in rats for 10 consecutive days beginning on the first postoperative day. Results were evaluated at three weeks postoperatively by measuring the sciatic functional index (SFI) at weekly intervals and the total number of nerve fibers and nerve fiber density of the nerve at three weeks. The SFI progressively improved for both irradiated and control nerves. Nerve fiber density also increased significantly in the irradiated nerves. They concluded that LLLT accelerates nerve regeneration. The difference between these results and the findings of the current study may be attributed to laser wave length they used being 904 nm in addition to daily irradiation.

Further study was carried out by Khullar et al.,⁹ who studied the effects of laser therapy on re-innervation of sciatic nerve after Gallium-Aluminum-Arsenide (GaAlAs) laser, 6J, 830 nm daily for a period of 28 days. They found improved functional recovery in the nerve indicated by SFI. They concluded that laser accelerated motor nerve re-innervation assessed by return of motor function subsequent to a standardized axonotmesis injury in the rat sciatic nerve. It can be noticed that the photo-chemical effects of LLL is greatly dose-dependent.

Recently, Zhang et al.,²⁰ conducted a study to discuss the effects of 660-nm GaAlAs LLL irradiation on neural regeneration after acellular nerve allograft repair of the sciatic nerve gap in rats. They found significant

increased nerve conduction velocity, restoration rate of tibialis anterior wet muscle weight and myelinated nerve number. They concluded that LLL therapy increased the rate of regeneration and target reinnervation after repair of the sciatic nerve. Consequently, LLL irradiation may be a useful, noninvasive adjunct for promoting nerve regeneration in surgically induced defects repaired nerves. Contrary to the results of the current study, their findings may be attributed to the difference in wave length (660 nm). Moreover, they applied laser for the sciatic nerve that may be more responsive to LLL than the facial one because of the anatomical pathway and the depth of penetration of laser.

There are several suggested mechanisms for acceleration of nerve regeneration. Tuner and Hode⁸ reported that LLLT helps the cell to produce ATP, the basic energy packer of cells- by up to 150% via affecting the mitochondria. This extra energy helps cells to promote healing. Laser light activates enzymes within the cells, causing them to produce proteins and rebuild themselves. Photons affect the cell membrane and normalize its polarity. This balanced polarity helps the cell to absorb the nutrient it needs and wastes more easily toxins from the cells.

Conclusion

Despite of the current easy availability of therapeutic laser and its trend to be routinely employed in injuries of a number of tissues, its effects on peripheral nerve regeneration are still controversial and need to be precisely determined before its application. According to the findings of the present research, it can be concluded that LLLT, with the parameters used in the current study, was not effective in improving functional outcome in BP.

REFERENCES

1. Anders, J.J., Borke, R.C., Woolery, S.K. and Van de Merwe, W.P.: Low power laser irradiation alters the rate of regeneration of the rat facial nerve. *Lasers Surg Med.* 13(1): 72-82, 1993.
2. Basford, J., Daube, J., Hallman, H., Millard, T. and Moyer, S.: Does Low-Intensity Helium-Neon Laser Irradiation Alter Sensory Nerve Active Potentials or Distal Latencies?. *Laser Surg Med.* 10: 35-39, 1991.
3. Beck, L. and Hall, J.: Evaluation of the Facial Nerve via Electroneuronography (ENoG). *The Hearing Journal*, 54(3): 36-45, 2001.
4. Chen, Y., Hsu, S., Chiu, C., Lin, J., Chen, C. and Yao, C.: Effect of Low-Power Pulsed Laser on Peripheral Nerve Regeneration in Rats. *Microsurgery*, 25: 83-89, 2005.
5. Endo, C., Barbieri, C., Mazzer, N. and Fasan, V.: Low-Power Laser Therapy Accelerates Peripheral Nerves' Regeneration. *Acta ortop Bras*, 16(5): 305-310, 2008.
6. Gilchrist, J.: Seventh Cranial Neuropathy. *Semin Neurol*, 29(1): 5-13, 2009.
7. Hato, N., Murakami, S. and Gyo, K.: Steroid and Antiviral Treatment for Bell's Palsy. *THE LANCET*, 371(9627): 1818-1820, 2008.
8. Kenyon, J.: *The Physiotherapist's Pocket Book: Essential facts at your fingertips.* Elsevier, China, 2008.
9. Khullar, S.: The Effect of Low Level Laser Treatment on Recovery of Conduction and Motor Function after Compression Injury in the Rat Sciatic Nerve. *Europ J Oral Sci*, 103: 299-305, 1995.
10. Kirshner, H.: *First Exposure to Neurology.* McGraw-Hill Inc., USA, 2007.
11. Lowe, A., Baxter, D., Walsh, D. and Allen, J.: Effect of Low Intensity Laser (830 nm) Irradiation on Skin Temperature and Antidromic Conduction Latencies in the Human Median Nerve: Relevance of radiant exposure. *Laser Surg Med*, 14: 40-46, 1994.
12. Ludman, H. and Bradley, J.: *ABC of Ear, Nose and Throat.* 5th edition, Blackwell Pub, Oxford, 2007.
13. May, M. and Schaitkin, B.: *The Facial Nerve.* 2nd edition, Thieme, New York, 2000.
14. Midamba, E. and Haanaes, H.: Low Reactive-Level 830 nm Ga Al As Diode Laser Therapy (LLLT) Successfully Accelerates Regeneration of Peripheral Nerves in Human. *Laser Therapy*, 5(3): 125-129.
15. Saitkulov, K., Shaimardanova, G. and Chelyshev, Y.: Regeneration of Nerve Fibers on Laser Irradiation of Spinal Nerve Projections. *Bulletin of Experimental Biology and Medicine*, 125: 309-311, 2006.
16. Tankéré, F. and Bernat, I.: Bell's Palsy: From viral etiology to diagnostic reality. *La Revue de Médecine Interne*, 30(9): 769-775, 2009.
17. Thomson, A., Skinner, A. and Piercy, J.: *Tidy's Physiotherapy.* 13th edition, Butterworth-Heinemann, Philadelphia, 2003.
18. Tuner, J. and Hode, L.: *The Laser Therapy Handbook.* 2nd edition, Prima books, Sweden, 2007.
19. Wong, I.: Surface Landmarks of the Facial Nerve Trunk: A prospective study. *ANZ J Surgery*, 71(12): 753-756, 2001.
20. Zhang, L., Tong, X., Yuan, X., Xiao-Hong Sun, X. and Jia, H.: Effects of 660-nm Gallium-Aluminum-Arsenide Low-Energy Laser on Nerve Regeneration after Acellular Nerve Allograft in Rats. *Synapse*, 64: 152-160, 2010.

الملخص العربي

تأثير أشعة الليزر ذو المستوى المنخفض على النتيجة الوظيفية في الأطفال المصابون بشلل بيل

الخلفية : عندما يصاب الشخص بشلل نصفي للوجه (شلل بيل) فإن استعادة الشكل الطبيعي ووظائف الوجه قد تستغرق مدة طويلة أو يحدث شفاء بشكل نسبي . **الهدف :** الهدف من هذه الدراسة هو تقييم فعالية العلاج بأشعة الليزر ذو المستوى المنخفض في الأطفال الذين يعانون من شلل بيل . **الأشخاص :** تم اختيار عشرين طفلاً مصابون بشلل بيل تراوحت أعمارهم من ١٢ إلى ١٥ عاماً و قُسموا عشوائياً إلى مجموعتين متساويتين (مجموعة دراسة و مجموعة ضابطة) . **الطريقة :** تم تقييم شدة إصابة العصب السابع باستخدام مقياس هاوس براكمان وكذلك مدى الاختلاف في الشكل الخارجي للوجه بالمقياس اليدوي باستخدام شريط القياس (المتر) . تلقت المجموعة الضابطة برنامج علاج طبيعي بالإضافة إلى أشعة الليزر الكاذبة بينما تلقت مجموعة الدراسة نفس البرنامج بالإضافة إلى العلاج بالليزر عن طريق تعريض عصب الوجه لأشعة الليزر تحت الحمراء المستمرة في بعض النقاط (مركز العصب السابع وبداية كل فرع من الفروع الحركية للعصب) بواقع أربع دقائق لكل نقطة بطول موجة ٨٣٠ نانو متر و قوة ١٠٠ ميلي وات (٦ جول/سم^٢) . استمر العلاج لمدة أربعة أسابيع بمعدل ثلاث جلسات أسبوعياً وكانت مدة الجلسة الواحدة ستون دقيقة . **النتائج :** أوضحت النتائج عدم وجود فروق ذات دلالة إحصائية بين المجموعتين في نتائج مقياس هاوس براكمان وتمائل نصفي الوجه . **الخلاصة :** يُستنتج من نتائج هذا البحث أن الليزر، بالمواصفات التي أُستخدمت في هذا البحث ، لم يكن له تأثير فعال في علاج الأطفال المصابون بشلل بيل .

الكلمات الدالة : مقياس هاوس براكمان - أشعة الليزر تحت الحمراء - شلل بيل .