

## Department of Biomechanics

### Doctoral Degree 2015

<b>Author</b>	:	<b>Hamada Ahmed Hamada Ahmed</b>
<b>Title</b>	:	<b>Mechanical Response of Hip and Knee Muscles Following Randomized Crossover Trials in Patellofemoral Pain Syndrome</b>
<b>Dept.</b>	:	<b>Department of Biomechanics.</b>
<b>Supervisors</b>	1.	<b>Salam Mohamed Elhafez</b>
	2.	<b>Ibrahim Ali Nassar</b>
	3.	<b>Hossam Eddien Fawaz</b>
<b>Degree</b>	:	<b>Doctoral.</b>
<b>Year</b>	:	<b>2015.</b>
<b>Abstract</b>	:	
<p><b>Background:</b> Faulty hip kinematics during weight bearing activities is proposed to contribute to patellofemoral pain syndrome (PFPS). However, limited information exists to determine the effectiveness of exercises programs that not only act on the knee joint, but also on the hip joint in these patients. <b>Purpose:</b> The purpose of this study was to compare between the effect of the starting of hip strengthening exercises before knee exercises program and the starting of knee exercises program before hip strengthening exercises on pain intensity, Kujala questionnaire scale, Q angle, anteversion angle, and knee extensors, hip abductors and lateral rotators peak torques (PT) in patients with PFPS. <b>Materials and Methods:</b> Twenty four patients suffering from PFPS were randomly assigned into two equal groups of twelve. Group (A) mean age, weight, height and BMI values were 23.33±5.39 years, 71.16±13.05 kg, 164.75±4.5 cm, and 26.21±4.71 kg/m<sup>2</sup> respectively. They received hip abductors and external rotators strengthening exercises for three weeks followed by knee extensors strengthening exercises and stretching exercises for quadriceps, hamstring, gastrocnemus muscles and iliotibial band for another three weeks. Group (B) with mean age, weight, height and BMI values were 23.16±6.33 years, 69.41±18.14 kg, 164.66±7.27 cm, and 25.2±6.2 kg/m<sup>2</sup> respectively. They received knee extensors strengthening exercises and stretching exercises for quadriceps, hamstring, gastrocnemus muscles, and iliotibial band for three weeks followed by hip abductors and external rotators strengthening exercises for another three weeks. Both groups received three to four sessions per week for six weeks. Pain level, Kujala scale, Q-angle, anteversion angle, and isokinetic eccentric PT for hip abductors, hip external rotators, and knee extensors were recorded before, after three weeks, and after six weeks of exercises. <b>Results:</b> 2x3 Mixed Design MANOVA revealed that there was a significant reduction in level of perceived pain and improvement of Kujala scale in group (A) compared with group (B) after six weeks of exercise (p&lt;0.05). However, there was no significant difference in the Q angle, anteversion angle, and isokinetic eccentric PT for hip abductors, hip external rotators and knee extensors between the tested groups after six weeks of exercise (p&gt;0.05). <b>Conclusion:</b> Starting rehabilitation program with hip strengthening exercises is more effective than starting with knee strengthening and stretching exercises in reducing pain and improving Kujala scale in patients with PFPS. Consequently, this may help physiotherapists in designing the most effective and efficient prevention and rehabilitation programs for patients suffering from PFPS.</p>		
<b>Key words</b>	1.	<b>Patellofemoral Pain Syndrome.</b>
	2.	<b>Hip, Knee.</b>
	3.	<b>Strengthening Exercises.</b>
	4.	<b>Q-Angle..</b>
	5.	<b>Anteversion Angle.</b>
	6.	
<b>Classification number</b>	:	<b>000.000.</b>
<b>Pagination</b>	:	<b>173 p.</b>
<b>Arabic Title Page</b>	:	<b>الاستجابة الميكانيكية لعضلات الفخذ والركبة عقب المحاولات العشوائية التبادلية لحالات آلام متلازمة الرضفة.</b>
<b>Library register number</b>	:	<b>4111-4112.</b>